

## Gulf of Mexico Harmful Algal Bloom Bulletin

Region: Texas


Monday, 06 May 2013

NOAA National Ocean Service

NOAA Satellite and Information Service

NOAA National Weather Service

Last bulletin: Monday, April 29, 2013



Satellite chlorophyll image with possible *K. brevis* HAB areas shown by red polygon(s), when applicable. Points represent cell concentration sampling data from April 26 to May 2: red (high), orange (medium), yellow (low b), brown (low a), blue (very low b), purple (very low a), pink (present), and green (not present). Cell count data are provided by Texas Parks and Wildlife Department. For a list of sample providers and a key to the cell concentration categories, please see the HAB-OFS bulletin guide:

[http://tidesandcurrents.noaa.gov/hab/habfs\\_bulletin\\_guide.pdf](http://tidesandcurrents.noaa.gov/hab/habfs_bulletin_guide.pdf)

Detailed sample information can be obtained through the Texas Parks and Wildlife Department at:

<http://www.tpwd.state.tx.us/landwater/water/enviroconcerns/hab/redtide/status.phtml>

To see previous bulletins and forecasts for other Harmful Algal Bloom Bulletin regions, visit the NOAA Harmful Algal Bloom Operational Forecast System bulletin archive:

<http://tidesandcurrents.noaa.gov/hab/bulletins.html>

## Conditions Report


There is currently no indication of a harmful algal bloom of *Karenia brevis* (commonly known as Texas red tide) at the coast in Texas. No respiratory impacts are expected alongshore the Texas coast today through Monday, May 13. For information on area shellfish restrictions, contact the Texas Department of State Health Services.

## Analysis

There is currently no indication of a harmful algal bloom of *Karenia brevis* at the coast in Texas. In MODIS Aqua imagery from 5/5 (shown left), patches of elevated chlorophyll (4 to 10  $\mu\text{g/L}$ ) are visible along- and offshore the Texas coastline from Sabine Pass to the Rio Grande. Patches of high to very high chlorophyll (10 to  $>20$   $\mu\text{g/L}$ ) are also visible along- and offshore from Sabine Pass to Port Aransas, including a patch of very high chlorophyll ( $>20$   $\mu\text{g/L}$ ) alongshore the southern end of Matagorda Island. Elevated chlorophyll is not indicative of the presence of *K. brevis* and is most likely due to the resuspension of benthic chlorophyll and sediments along the coast.

Forecast models based on predicted near-surface currents indicate a potential maximum transport of 40 km north from the Port Aransas region from May 5-9.


Derner, Davis



Wind speed and direction are averaged over 12 hours from buoy measurements. Length of line indicates speed; angle indicates direction. Red indicates that the wind direction favors upwelling near the coast. Values to the left of the dotted vertical line are measured values; values to the right are forecasts. Wind observation and forecast data provided by NOAA's National Weather Service (NWS).


## Wind Analysis

**Port Aransas:** Southeast winds (5-15kn, 3-8m/s) today shifting south after midnight. South winds (5-10kn, 3-5m/s) Tuesday becoming southeast (10-15kn, 5-8m/s) Tuesday afternoon through Thursday, increasing to 15-20kn (8-10m/s) Friday. South winds (10-15kn) Friday night.



Satellite chlorophyll image and forecast winds for May 7, 2013 06Z with points representing cell concentration sampling data from April 26 to May 2: red (high), orange (medium), yellow (low b), brown (low a), blue (very low b), purple (very low a), pink (present), and green (not present). Cell count data are provided by Texas Parks and Wildlife Department. For a list of sample providers and a key to the cell concentration categories, please see the HAB-OFS bulletin guide:

[http://tidesandcurrents.noaa.gov/hab/habfs\\_bulletin\\_guide.pdf](http://tidesandcurrents.noaa.gov/hab/habfs_bulletin_guide.pdf)



Verified and suspected HAB areas shown in red. Other areas of high chlorophyll concentration shown in yellow (see p. 1 analysis for interpretation).